

No Spontaneous CP Violation at Finite Temperature in the MSSM?^a

P. John

*Instituto de Estructura de la Materia (CSIC), Serrano 123, E-28006 Madrid, Spain
E-mail: john@makoki.iem.csic.es***Abstract**

In order to generate the baryon asymmetry of the Universe sufficiently strong CP violation is needed. It was therefore proposed that at finite temperature there might be spontaneous (transitional) CP violation within the bubble walls at the electroweak phase transition in supersymmetric models. We investigate this question in the MSSM.

1 Introduction

For producing the baryon asymmetry of the Universe, we need extensions to the Standard Model. One of the Zakharov conditions requires nonequilibrium. In the MSSM this can be fulfilled by a strong enough first order phase transition with a light scalar top¹. Also a small bubble wall velocity² seems to support baryogenesis. But Zakharov's conditions also require CP violation and in the MSSM there are several mechanisms known to generate it. Explicit CP violating operators might conflict to experimental EDM bounds³. It were interesting to have a mechanism generating enough CP violation for baryogenesis without any conflict to experiments. While spontaneous CP violation is excluded at $T = 0$ for the experimentally allowed parameter values⁴, there is a suggestion that it might be more easily realized at finite temperatures^{5,6}.

Previously, the moduli of the two Higgs doublets around the phase boundary have been determined from the 2-loop effective potential^{7,8}. The CP violating phase between the two Higgs doublets has been addressed perturbatively^{9,10} and nonperturbatively¹¹.

We present the first complete solution of the equations of motion for the phase between the two Higgs doublets within the MSSM, utilizing a perturbative effective potential, but without restricting it to the effective quartic couplings. Our conclusions¹⁰ differ from those obtained earlier on.

2 Searching for CP violating phases

We parameterize the two Higgs doublets of the MSSM as

$$H_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} h_1 e^{i\theta_1} \\ 0 \end{pmatrix}, \quad H_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ h_2 e^{i\theta_2} \end{pmatrix}. \quad (1)$$

In addition, because of gauge invariance, the effective Higgs potential depends on the phases only via $\theta = \theta_1 + \theta_2$, and we have an additional constraint $h_1^2 \partial_\mu \theta_1 = h_2^2 \partial_\mu \theta_2$. We can then concentrate on θ . Assuming tree-level kinetic terms and

^aTo appear in the Proceedings of SEWM2000, Marseille, June 14-17, 2000

moving to a frame where the bubble wall is static and planar, the action to be minimized is

$$S \propto \int dz \left[\frac{1}{2}(\partial_z h_1)^2 + \frac{1}{2}(\partial_z h_2)^2 + \frac{1}{2} \frac{h_1^2 h_2^2}{h_1^2 + h_2^2} (\partial_z \theta)^2 + V_T(h_1, h_2, \theta) \right], \quad (2)$$

where $V_T(h_1, h_2, \theta)$ is the finite temperature effective potential for h_1, h_2, θ . In general, we are solving the equations of motion for h_1, h_2, θ following from this action. In the numerical solution we use the method outlined in ⁸ which deals with the minimization of a functional of the squared equations of motion.

At the first stage, we consider the case with no explicit CP phases, and ask whether a particular solution without CP violation ($\theta = 0, \pi$), is in fact a local minimum of the action or not. Clearly, it is not if

$$m_3^2(h_1, h_2) \equiv \frac{1}{|h_1 h_2|} \left. \frac{\partial^2 V_T(h_1, h_2, \theta)}{\partial \theta^2} \right|_{\theta=0} < 0, \quad (3)$$

where we have divided by $|h_1 h_2|$, assuming that this is non-zero. Eq. (3) is to be evaluated along the path found by solving the equations of motion for h_1, h_2 . We have chosen the convention that h_1 can have either sign, allowing us to consider only $\theta = 0$. For the case of the most general quartic two Higgs doublet potential, Eq. (3) agrees with the constraint on which most of the investigations of spontaneous CP violation are based. However, Eq. (3) is true more generally, independent of the form of the potential $V_T(h_1, h_2, \theta)$.

The tree-level potential of the theory is

$$V_{\text{tree}} = \frac{1}{2} m_1^2 h_1^2 + \frac{1}{2} m_2^2 h_2^2 + m_{12}^2 h_1 h_2 \cos \theta + \frac{1}{32} (g^2 + g'^2) (h_1^2 - h_2^2)^2, \quad (4)$$

where g, g' are the SU(2) and U(1) gauge couplings, and at tree-level

$$m_{12}^2 = -\frac{1}{2} m_A^2 \sin 2\beta. \quad (5)$$

It follows that $m_3^2(h_1, h_2) = (1/2) m_A^2 \sin 2\beta > 0$, so that the minimum of the potential in the θ direction is at $\theta = 0$. Thus, in order to get spontaneous CP violation one needs radiative corrections which can overcome the tree-level term.

Older considerations^{5,6,9} are based on the approximation to the effective potential where only the quadratic and quartic operators are considered. At finite temperatures around the electroweak phase transition, important contributions come from infrared sensitive non-analytic contributions which are not of this form, and can affect spontaneous CP violation^{11,10}. Thus, it is important to solve the equations of motion more generally for the full effective potential.

Here we consider the full finite temperature 1-loop effective potential of the MSSM. It is known that 2-loop corrections are very important in the MSSM¹, allowing for larger values of h_1, h_2 in the broken phase. Nevertheless, for the present problem we find that even 1-loop effects are in most cases very small, so we do not expect qualitative changes from the 2-loop effects.

3 A scan for spontaneous transitional CP violation.

The tree-level part of the effective potential $V_T(h_1, h_2, \theta)$ is in Eq. (4). In the resummed 1-loop contribution to $V_T(h_1, h_2, \theta)$, we include gauge bosons, stops, charginos and neutralinos. This introduces dependences on the trilinear squark

mixing parameters A_t and μ as well as on the squark mass parameters m_Q^2 , m_U^2 , and the U(1), SU(2) gaugino parameters M_1 and M_2 .

We now wish to see whether the constraint in Eq. (3) can be satisfied at the bubble wall between the symmetric and broken phases. To do so, we have to search for each parameter set for the critical temperature T_c , solve the equations of motion for (h_1, h_2) between the minima, and evaluate $m_3^2(h_1, h_2)$ along this path. Since this is quite time-consuming, we proceed in two steps.

1. At the first stage, we do not solve for h_1, h_2, T_c , but rather take them as free parameters in the ranges $h_1/T = -2..2$ and $h_2/T = 0..2$, $T = 80..120$ GeV. The zero temperature parameters are varied in the wide ranges

$$\begin{aligned} \tan \beta &= 2..20, & m_A &= 0..400 \text{ GeV}, \\ m_U &= -50..800 \text{ GeV}, & m_Q &= 50..800 \text{ GeV}, \\ \mu, A_t, M_1, M_2 &= -800..800 \text{ GeV}. \end{aligned} \tag{6}$$

Here a negative m_U means in fact a negative right-handed stop mass parameter, $-|m_U^2|$. We have also studied separately the (dangerous¹²) region where the transition is very strong^{1,12}, corresponding to $m_U \sim -70... -50$ GeV.

Note that since we do not solve for the equations of motion at this stage but allow for $h_1 = \pm|h_1|$, we have to divide in Eq. (3) by $h_1 h_2$ instead of $|h_1 h_2|$: this leads in general to positive values due to the tree-level form of the potential, Eqs. (4),(5). A signal of a potentially promising region is then a small absolute value of the result, since this means that we are close to a point where $\partial_\theta^2 V_T(h_1, h_2, \theta)$ crosses zero.

2. At the second stage, we study the most favourable parameter region thus found in more detail. First of all, we search for the critical temperature. Then, we solve the equations of motion for (h_1, h_2) . By comparing with the exact numerical solution in several cases, we find that a sufficient accuracy can be obtained in practice by searching for the “ridge” as an approximation to the wall profile. It is determined as the line of maxima of the potential in the direction perpendicular to the straight line between the minima. Finally, we look for the minimum of $V_T(h_1, h_2, \theta)$ at fixed (h_1, h_2) : this is a fast and reliable approximation for the full solution in the case that θ is small (i.e., just starts to deviate from zero), and corresponds to Eq. (3).

For the first stage, we perform a Monte Carlo scan with about $2 \cdot 10^9$ configurations. Small values of $m_3^2(h_1, h_2)$ are scarce, and even then do not necessarily correspond to the desired phenomenon of spontaneous CP violation: they could also be points far from the actual wall. This can be clarified at stage 2.

The parameter region found depends most strongly on m_A , $\tan\beta$, with a preference on small values of m_A and large of $\tan\beta$, such that m_{12}^2 in Eq. (5) is small. (This is in contrast to the requirements of a strong phase transition¹). There is also a relatively strong dependence on A_t and μ : the region favoured is shown in Fig. 1. The dependences on the other parameters are less significant; for m_U and m_Q small values are preferred. The region found is in rough agreement with those found in ^{5,6,9,11}.

At the second stage, we make further restrictions. For instance, we exclude the cases leading to non-physical negative mass parameters. We also exclude cases leading to $T = 0$ spontaneous CP violation in the broken phase: this phenomenon requires very small values⁴ of m_A . We also discard phase transitions which are exceedingly weak, $v/T \ll 0.1$.




Figure 1: The average value of m_3^2 versus μ and A_t . We observe that small values of m_3^2 are not typical in any part of the plane but are on the average more likely for small μ, A_t , and that the distribution is wider (and thus more favourable) for like signs of μ, A_t , as shown by the noisy contours obtained with a finite amount of statistics.

In⁹ the special point $m_U^2 \approx 0$ was considered. Since in⁹ the thermal mass corrections were neglected this corresponds in the physical MSSM to a case where $m_U^2 + \#T^2 \sim 0$. Expanding the 1-loop cubic term from the stops to a finite order in v_1/v_2 , it was suggested that transitional spontaneous CP violation can take place. This region is quite dangerous due to the vicinity of a charge and colour breaking minimum¹². Without expanding the 1-loop contribution in v_1/v_2 , we cannot reproduce the behaviour proposed there. In any case, even before taking into account the experimental lower limits on the Higgs masses, we cannot find any promising case in the sample of $\sim 2 \times 10^6$ configurations of stage 2.

We conclude¹⁰ that after taking into account the infrared sensitive effects inherent in the 1-loop effective potential, coming from a light stop and gauge bosons, and solving for the wall profile from the equations of motion, spontaneous CP violation does not take place in the physical MSSM bubble wall.

Acknowledgements I would like to thank S. Huber, M. Laine, M. Schmidt for collaboration in¹⁰.

1. J.R. Espinosa, *Nucl. Phys.* **B475** (1996) 273; Bödeker, John, Laine, Schmidt, *Nucl. Phys.* **B497** (1997) 387; Carena, Quirós, Wagner, *Nucl. Phys.* **B524** (1998) 3; Losada, hep-ph/9905441; Laine, Rummukainen, *Nucl. Phys.* **B535** (1998) 423; *ibid.*, *Phys. Rev. Lett.* **80** (1998) 5259.
2. G. Moore, JHEP 0003:006,2000; P. John and M.G. Schmidt, hep-ph/0002050, Proc of *SEWM 2000*.
3. Discussion and refs.: Pilaftsis and Wagner, *Nucl. Phys. B* 553 (1999) 3.
4. N. Maekawa, *Phys. Lett. B* 282 (1992) 387; A. Pomarol, *Phys. Lett. B* 287 (1992) 331.
5. D. Comelli and M. Pietroni, *Phys. Lett. B* 306 (1993) 67; J.R. Espinosa, J.M. Moreno and M. Quirós, *Phys. Lett. B* 319 (1993) 505.
6. D. Comelli, M. Pietroni and A. Riotto, *Nucl. Phys. B* 412 (1994) 441.
7. J.M. Moreno, M. Quirós, M. Seco, *Nucl. Phys.* **B526** (1998) 489; J. Cline, G. Moore, *Phys. Rev. Lett.* **81** (1998) 3315.
8. P. John, *Phys. Lett.* **B452** (1999) 221; P. John, Proc. of *SEWM 98*.
9. K. Funakubo, A. Otsuki, F. Toyoda, *Prog. Theor. Phys.* 102(1999) 389.
10. S.J. Huber, P. John, M. Laine, M. Schmidt, *Phys. Lett. B* 475 (2000) 104.
11. M. Laine and K. Rummukainen, *Nucl. Phys. B* 545 (1999) 141, Proc. of *Lattice '99*, hep-lat/0009025; Z. Fodor, Proc. of *Lattice '99*.
12. J.M. Cline, G.D. Moore and G. Servant, *Phys. Rev. D* 60 (1999) 105035.